

## Phenolic acid content in wheat grain (*Triticum* spp) of different genotypes

### Contenido de ácido fenólico (*Triticum* spp) de diferentes granos de trigo

Danuta Kurasiak-Popowska <sup>1</sup>, Kinga Stuper-Szablewska <sup>2</sup>,  
Jerzy Nawracała <sup>1</sup>, Agnieszka Tomkowiak <sup>1</sup>, Juliusz Perkowski <sup>2</sup>

Originales: *Recepción*: 06/02/2015- *Aceptación*: 14/12/2015

*Nota científica*

#### ABSTRACT

One of the most significant factors affecting resistance to fungal diseases in winter wheat genotypes is the content of phenolic compounds. A total of 100 winter wheat cultivars were investigated. The contents of five phenolic acids: ferulic, vanillic, syringic, vanillin, p-coumaric and free phenolic acids (FPA) were determined. Considerable variation was found both within tested cultivars. Among the investigated acids the greatest amounts were recorded for ferulic acid and its mean concentration was 975  $\mu\text{g g}^{-1}$ , p-coumaric acid was characterised by a lower mean concentration (52  $\mu\text{g g}^{-1}$ ), while the lowest concentration was recorded for vanillin, amounting to 5  $\mu\text{g g}^{-1}$ . Ferulic acid state 92.4% of total phenolic acids. Significant differences in content of ferulic acids between genotypes were observed. The group of winter wheats coming from different European countries was characterized by high variation in the contents of ferulic acid, ranging from 320 to 3103  $\mu\text{g g}^{-1}$ . No significant correlations were observed between ferulic acid and other phenolic acids. Results provide the basis for further studies in this field.

#### Keywords

ferulic acid • phenolic acids • wheat

- 
- 1 Department of Genetics and Plant Breeding. Faculty of Agronomy and Bioengineering. Poznan University of Life Sciences. Dojazd 11, 60-632 Poznań, Poland. popowska@up.poznan.pl
  - 2 Department of Chemistry. Faculty of Wood Technology. Poznan University of Life Sciences. Wojska Polskiego 75, 60-625 Poznań, Poland.

## RESUMEN

Uno de los factores más importantes en la resistencia a las enfermedades fúngicas en genotipos de trigo de invierno es el contenido de compuestos fenólicos. Se investigó un total de 100 cultivos de trigo de invierno, se determinó sus contenidos de los cinco ácidos fenólicos: ferúlico, vanílico, siríngico, vainillina, p-cumárico y ácidos fenólicos libres (AFL) y se encontró una variación entre ambos cultivos probados. Entre los ácidos investigados, se detectó mayores cantidades de ácido ferúlico (concentración media de  $975 \mu\text{g g}^{-1}$ ), ácido p-cumárico (caracterizado por una baja concentración media ( $52 \mu\text{g g}^{-1}$ ), mientras que la concentración más baja que se detectó fue de vainillina (a razón de  $5 \mu\text{g g}^{-1}$ ). El ácido ferúlico conforma el 92,4% de los ácidos fenólicos totales. Por otro lado, se observó diferencias significativas en el contenido de ácidos ferúlicos entre genotipos: el grupo de trigos de invierno procedente de varios países europeos tenía como característica una alta variación de contenidos de ácido ferúlico, del orden de 320 hasta  $3103 \mu\text{g g}^{-1}$ . No se detectó correlaciones significativas entre el ácido ferúlico y otros ácidos fenólicos. En conclusión, los resultados de este estudio ofrecen la base para otras investigaciones con más profundidad en este ámbito.

### Palabras clave

ácido ferúlico • ácidos fenólicos • trigo

## INTRODUCTION

Wheat is infested by many fungal diseases. In case of wheat this problem to the greatest extent pertains to grain contamination with mycotoxins produced by various species, of which the most important are fungi from the genus *Fusarium* (2, 15). This results in grain contamination with such toxins as deoxynivalenon (DON), nivalenol (NIV) (14).

Breeding towards resistance is the best prevention method. Although some genotypes are known to have resistance coming from spring cultivars such as Sumai 3 and Frontana, they exhibit disadvantageous yield traits. In view of such difficulty in obtaining resistant cultivars to *Fusarium* new genotypes are searched for both in Poland and worldwide, which could provide good germplasm for breeding towards resistance.

Some information indicates that also resistance to Fusarium Head Blight (FHB) in wheat may be to a considerable degree connected with the presence of phenolic acids in the spike and in kernels. Among phenolic acids in wheat grain ferulic acid is found in greatest amounts, which level depends on the cultivar and on kernel size (8, 10).

A dependence was indicated between ferulic acid found in wheat grain and resistance of wheat to fusariosis (17).

The aim of the presented investigations was to determine the contents of phenolic acids, particularly ferulic acid, in grain of winter wheat genotypes.

## MATERIAL AND METHODS

A total of 100 cultivars and lines of winter wheat were selected for analysis. They came from Polish breeding companies involved in wheat breeding and from the collection of winter wheat belonging to the Department of Genetics and Plant Breeding, the Poznań University of Life Sciences. Breeding companies, *i.e.* Małopolska Plant Growing Company - HBP Ltd. (MHR), Plant Breeding Company Smolice Ltd. (HRS), Danko Plant Breeding Company Ltd. (Danko HR), Poznań Plant Breeders Ltd., (PHR), Plant Breeding Company Strzelce Ltd. (HRSt) from their own breeding material selected for analyses genotypes with known resistance to fungal diseases. Genotypes selected from the collection of the Department of Genetics and Plant Breeding, PULS included cultivars and lines of winter wheat originated from different countries (Austria, Czech Republic, Holland, France, Germany, Russia and Hungary).

Based on literature data they were considered resistant to *Fusarium*, and they were obtained from the wheat collection of the National Small Grain Collection at the Agriculture Research Station in Aberdeen, belonging to the National Plant Germplasm System in the USA, and from dr Tomasz Góral, Plant Breeding and Acclimatization Institute in Radzików. Analyses were also conducted on wheat lines generated at the Department of Genetics and Plant Breeding (DG&PB) and coming from crossing of semi-dwarf English and French forms with leading Polish cultivars.

All grain samples used in the analyses were harvested in 2011 from genotypes growing at Wielkopolska conditions, at Agriculture Research Station "Dłóż".

The soil at the site is Luvisols (SgP, 2011) with pH ranged between 6.0 and 6.4. Weather conditions in the spring/vegetation in 2011 were as follows: the average temperature in April was 11.9°C, 15.5°C in May, June 19.0°C and in July 18.6°C. Total rainfall in the month of April was 14 mm, 37.5mm in May, 33.5mm in June and 169mm in July. The field experiment was conducted in 3 replications, while the plot size was 1m<sup>2</sup>.

From 3 replications, a bulk sample of 1 kg was collected, from which a sample of 100 g grain each was used in further analyses. The chemical analysis were conducted in 2 replication, and the mean of the replications is shown in the tables (table 1, page 4)

### 2.1 Analysis of contents of free phenolic acids (FPA)

Total of free phenolic acids was determined as described by Stuper-Szablewska *et al.*, (2015).

### 2.2 Determination of phenolic acids

Contents of 5 phenolic acids was determined as described by Stuper-Szablewska *et al.*, (2014).

### 2.3 Statistical analysis

Results were subjected to statistical analysis using the STATISTICA v. 8.0 software package. Means, range from the samples as well as the number and percentage of positive samples were determined. Moreover, values of Pearson's linear correlation coefficients were determined at the following significance levels,  $\alpha=0.01$ , and  $\alpha=0.001$  (\*\*, \*\*\*). In order to compare contents of analysed metabolites in samples the procedure of multiple comparisons using Tukey's method was applied.

**Table 1.** Mean, ranges and percentages of positive samples for concentrations of five identified phenolic acids and their totals as well as free phenolic acids (FPA).

**Tabla 1.** Media, rangos y porcentajes de muestras positivas para las concentraciones de cinco ácidos fenólicos identificados, sus totales y ácidos fenólicos libres (AFL).

	Ferulic $\mu\text{g g}^{-1}$	Vanillic $\mu\text{g g}^{-1}$	Syringic $\mu\text{g g}^{-1}$	Vanillin $\mu\text{g g}^{-1}$	p-cumaric $\mu\text{g g}^{-1}$	Phenolic acids (total) $\mu\text{g g}^{-1}$	FPA $\mu\text{g g}^{-1}$
Mean	975	17	5	6	52	1055	515
Range	320-3103	0-72	0-37	0-77	10-160	357-3186	220-843
Percent of positive samples	100%	71%	40%	38%	100%	100%	100%

**Table 2.** Co-occurrence of phenolic acids.

**Tabla 2.** Co-ocurrencia de ácidos fenólicos.

Co-occurrence	Percent of positive samples
Ferulic, vanillic, syringic, vanillin, p-cumaric	11
Ferulic, vanillic, vanillin, p-cumaric	23
Ferulic, vanillic, syringic, p-cumaric	14
Ferulic, syringic, vanillin, p-cumaric	3
Ferulic, vanillin, p-cumaric	3
Ferulic, syringic, p-cumaric	9
Ferulic, vanillic, p-cumaric	19
Ferulic, p-cumaric	18

## RESULTS

A total of 100 grain samples of winter wheat were analysed. In the samples five phenolic acids were identified: ferulic, vanillic, syringic, vanillin and p-coumaric acids (table 1).

However, all the acids were presented in grain of only 11 genotypes (table 2).

Most frequently in the analysed samples (23%) four acids were found simultaneously: ferulic, vanillic, vanillin and p-coumaric acids as well as a combination of three acids: ferulic, vanillic and p-coumaric (table 2). In grain of all the genotypes two acids were detected: ferulic and p-coumaric acids.

Mean concentration of ferulic acid detected in 100% samples was  $975 \mu\text{g g}^{-1}$  while that of p-coumaric acid - also found in all samples - was  $52 \mu\text{g g}^{-1}$ , with a considerable range from the sample observed in both cases. The presence of vanillin was found in grain of 38% tested genotypes, while syringic acid was found in grain of 40%.

At the same time in both cases the lowest mean content of these acids was recorded, amounting to 6 and  $5 \mu\text{g g}^{-1}$ , respectively.

Calculated linear correlation coefficients showed that contents of ferulic acid and total phenolic acids were related to the highest degree, highly significantly (table 3, pág. 5).

**Table 3.** Matrix of internal correlations for concentrations of free phenolic acids (FPA), five identified phenolic acids and their totals.**Tabla 3.** Matriz de correlaciones internas para las concentraciones de ácidos fenólicos libres (AFL), cinco ácidos fenólicos identificados y sus totales.

	FPA	Ferulic	Vanilic	Syringic	Vanilin	p-cumaric
Ferulic	0.4801**					
Vanilic	-0.0238	0.0732				
Syringic	0.4193**	0.1205	-0.1679			
Vanilin	0.0232	-0.0085	0.0551	-0.0907		
p-cumaric	-0.0347	-0.0284	0.0036	0.0426	-0.0301	
phenolic acids (total)	0.4824**	0.9974***	0.1065	0.1315	0.0129	0.0291

\*\* ,\*\*\* - Pearson's linear correlation coefficients at the significance level  $\alpha = 0.01$ ,  $\alpha = 0.001$ .

\*\* ,\*\*\* - coeficientes de correlación lineal de Pearson en el nivel de significación  $\alpha = 0,01$ ,  $\alpha = 0,001$ .

This dependence resulted from the fact that ferulic acid constituted 92.4% total phenolic acids. Significant correlations were also recorded between contents of ferulic acid, syringic acid and total phenolic acids and contents of FPA (table 3).

Conducted analyses of winter wheat grain showed very high variation in the analysed genotypes in terms of contents of ferulic acid (320-3103  $\mu\text{g g}^{-1}$ ) and total phenolic acids and FPA (table 1, page 4; table 2, page 4; table 3). Almost half of the genotypes: 47, had contents of ferulic acid within the range of 700-1000  $\mu\text{g}$ , and only thirteen contained over 1500  $\mu\text{g}$ . A very low content of ferulic acid <500  $\mu\text{g}$  was found in only five genotypes.

## DISCUSSION

Li *et al.* (2008) when analysing 130 genotypes of winter wheat found that the range from samples for ferulic acid was from 326 to 1171  $\mu\text{g g}^{-1}$  at the mean contents of 664  $\mu\text{g g}^{-1}$ , while in this experiment the average for winter wheat was 974  $\mu\text{g g}^{-1}$ . Significant differences between contemporary and old cultivars

of *Triticum aestivum* and *T. durum* were reported by Heimler *et al.* (2010). Also the analysis of 58 genotypes from the Australian Winter Cereals Collection conducted by Wu *et al.* (2001) showed significant differences between genotypes in terms of contents of seven phenolic acids. Irmak *et al.* (2008) stated that even in isolines of cv. 'Pegaso' differing in glutenin protein subunits there were significant differences in PFA contents and total phenolic acids. Probably reports on a lack of significant differences between genotypes frequently result from a small number of analysed forms (3, 21). Amounts of ferulic, coumaric and vanillic acids comparable to those recorded in these experiment were observed in different wheat cultivars by other authors (11, 12).

Analogously to different publications on wheat phenolics, ferulic acid was the dominant phenolic acid in all wheat cultivars and accounted for 92.4% of total phenolic acids. Similarly, Irmak *et al.* (2008) and Jonnalala *et al.* (2010) showed that ferulic acid constituted 92% and 95%, respectively, of total phenolic acids. Moreover, a high correlation was

found between contents of ferulic acid and total contents of phenolic acids (12).

Significant differences in contents of ferulic acid and total contents of phenolic acids between spring wheat genotypes were found by Mpofo *et al.* (2006). Content of ferulic acid was low, amounting to 371-436  $\mu\text{mol gallic acid } 100\text{g}^{-1}$  grain, while the total content of phenolic acids was high, amounting to 1709-1990  $\mu\text{g g}^{-1}$ . In turn, Irmak *et al.* (2008) recorded that total phenolic acid content in cv. 'Pegas' was very high at 2795  $\mu\text{g g}^{-1}$  GAE. Kim *et al.* (2006) in wheat samples recorded concentrations of ferulic acid at an average level of 3420  $\mu\text{g g}^{-1}$ .

Content of FPA was on average over two-fold lower than total phenolic acids.

Greater differences were observed by Okarter *et al.* (2010), as in their study FPA content ranged from 255 to 499  $\mu\text{mol gallic acid } 100\text{g}^{-1}$  grain, while total content of phenolic acids from 841 to 1099  $\mu\text{mol gallic acid } 100\text{g}^{-1}$  grain, with significant differences between genotypes. In the analyses performed on 11 cultivars by Adom *et al.* (2003), FPA content was much lower (119-201  $\mu\text{mol gallic acid } 100 \text{ g}^{-1}$  grain) than the total content of phenolic acids at 709-860  $\mu\text{mol gallic acid } 100 \text{ g}^{-1}$  grain.

Considerable differences in contents of ferulic acid, observed in the conducted experiment, may be reflected in differences in resistance to *Fusarium*.

## REFERENCES

1. Adom, K. K.; Sorrells, M. E.; Liu, R. H. 2003. Phytochemical profiles and antioxidant activity of wheat varieties. *Journal of Agricultural and Food Chemistry*. 51: 7825-7834.
2. Chiotta, M. L.; Chulze, S.; Barros, G. 2015. Fuentes de inóculo de especies de *Fusarium* potenciales productoras de micotoxinas en el agroecosistema soja. *Revista de la Facultad de Ciencias Agrarias. Universidad Nacional de Cuyo. Mendoza. Argentina*. 47(2): 171-184.
3. Gélinas, P.; McKinnon, C. M. 2006. Effect of wheat variety, farming site, and bread baking on total phenolics. *International Journal of Food Science and Technology*. 41: 329-332.
4. Heimler, D.; Vignolini, P.; Isolani, L.; Arfaioli, P.; Ghiselli, L.; Romani, A. 2010. Polyphenol content of modern and old varieties of *Triticum aestivum* L. and *T. durum* Desf. grains in two years of production. *Journal of Agricultural and Food Chemistry*. 58: 7329-7334.
5. Irmak, S.; Jonnala, R. S.; MacRitchie, F. 2008. Effect of genetic variation on phenolic acid and policosanol contents of Pegaso wheat lines. *Journal of Cereal Science*. 48: 20-26.
6. Jonnala, R. S.; Irmaka, S.; MacRitchie, F.; Bean, S. R. 2010. Phenolics in the bran of waxy wheat and triticale lines. *Journal of Cereal Science*. 52: 509-515.
7. Kim, K. H.; Tsao, R.; Yang, R.; Cui, S. W. 2006. Phenolic acid profiles of wheat extracts and the effect of hydrolysis conditions. *Food Chemistry*. 90: 426-433.
8. Lempereur, I.; Rouau, X.; Abecassis, J. 1997. Genetic and agronomic variation in arabinoxylan and ferulic acid contents of durum wheat (*Triticum durum* L.) grain and its milling fractions. *Journal of Cereal Science*. 25: 103-110.
9. Li L.; Shewry, P. P.; Ward, J. L. 2008. Phenolic acids in wheat varieties in the HEALTH GRAIN diversity screen. *Journal of Agricultural and Food*. 56: 9732-9739.
10. Manfreda, V. T.; Acosta, M. C. 2015. Variables de tamaño y forma de granos y embriones de trigo (*Triticum aestivum* L.): análisis y propuestas. *Revista de la Facultad de Ciencias Agrarias. Universidad Nacional de Cuyo. Mendoza. Argentina*. 47(1): 43-58.
11. Moore, J.; Hao, Z.; Zhou, K.; Luther, M.; Costa, J.; Yu L. 2005. Carotenoid, tocopherol, phenolic acid and antioxidant properties of Maryland-grown softwheat. *Journal of Agricultural and Food Chemistry*. 53: 6649-6657.

12. Mpofu, A.; Sapirstain, H. D.; Beta, T. 2006. Genotype and environmental variation in phenolic content, phenolic acid composition, and antioxidant activity of hard spring wheat. *Journal of Agricultural and Food Chemistry*. 54: 1265-1270.
13. Okarter, N.; Liu, C.; Sorrells, M. E.; Liu, R. H. 2010. Phytochemical content and antioxidant activity of six diverse varieties of whole wheat. *Food Chemistry*. 119: 249-257.
14. Perkowski, J.; Wiwart, M.; Buśko, M.; Laskowska, M.; Berthiller, A.; Kandler, S.; Karska, R. 2007. Fusarium toxins and total fungal biomass indicators in naturally contaminated wheat samples from north-eastern Poland in 2003. *Food Additives and Contaminants*. 24: 1292-1298.
15. Perniola, O. S.; Staltari, S.; Chorzempa, S. E.; Astiz Gassó, M. M.; Molina, M. del C. 2014. Control biológico de *Fusarium graminearum*: utilización de *Trichoderma* spp. y biofumigación con parte aérea de *Brassica juncea*. *Revista de la Facultad de Ciencias Agrarias. Universidad Nacional de Cuyo. Mendoza. Argentina*. 46(2): 45-56.
16. SgP. 2011. Systematyka gleb Polski. *Soil Science Annual (Roczniki Gleboznawcze)*. 62(3):5-142.
17. Stuper K.; Perkowski, J.; Nawracała, J. 2011. Comparison of analytical methods of phenolic compounds determined in cereal grains. *New trends in food analytics. M Szoltyś i A Dąbrowska (eds.)*. p. 107-118.
18. Stuper-Szablewska, K.; Kurasiak-Popowska, D.; Nawracała, J.; Perkowski, J. 2014. Comparison of phenolic acids contents in various wheat genotypes. *Przemysł Chemiczny*. 93/12: 2274-2278.
19. Stuper-Szablewska, K.; Ostrowska, A.; Góral, T.; Matysiak, A.; Perkowski, J. 2015. Comparison of the antioxidant activity of different varieties of winter wheat naturally infected and inoculated with fungi of the genus *Fusarium*, *Biuletyn IHAR*. 276: 9-18.
20. Wu, H.; Haig, T.; Pratley, J.; Lemerle, D.; An, M. 2001. Allelochemicals in wheat (*Triticum aestivum* L.): variation of phenolic acids in shoot tissues. *Journal of Chemical Ecology*. 27: 125-135.
21. Zhou, K.; Laux, J. J.; Yu, L. 2004. Comparison of Swiss red wheat grain and fractions for their antioxidant properties. *Journal of Agricultural and Food Chemistry*. 52: 1118-1123.